


PEPTIDE AND PROTEIN RADIOLABELING

Fig. 1. Radiosynthesis and HPLC purification of [¹²⁵I]iodo-thyronine prepared using the iodogen method.




HPLC conditions

Column: C18 4.7 x 150 mm

Mobile phase: Gradient 25 – 50 % ACn, 0.1 % TFA

Flow rate: 1 ml/min

Fig. 2. Radiosynthesis and HPLC purification of [¹²⁵I]iodo-[D-Ala²]deltorphan II prepared via the lactoperoxidase method.




HPLC conditions

Column: C18 4.7 x 150 mm

Mobile phase: Gradient 25 – 60 % ACn, 0.1 % TFA

Flow rate: 1 ml/min

Fig 3. Radiosynthesis and HPLC purification of [¹²⁵I]iodo-SHU 9119 prepared via the lactoperoxidase method.




HPLC conditions

Column: C18 4.7 x 150 mm

Mobile phase: Gradient 25 – 50 % ACn, 0.1 % TFA

Flow rate: 1 ml/min

Fig. 4. Radiosynthesis and HPLC purification of [¹²⁵I]iodo-GIP prepared using a soluble oxidizing agent (N-chlorosuccinimide).




HPLC conditions

Column: C18 4.7 x 50 mm

Mobile phase: Gradient 25 – 50 % ACn, 0.1 % TFA

Flow rate: 1 ml/min

Fig. 5. QC check on HPLC-purified [¹²⁵I]iodo-orexin.



HPLC conditions

Column: C18 4.7 x 150 mm

Mobile phase: Gradient 25 – 40 % ACn, 0.1 % TFA

Flow rate: 1 ml/min